

## **Thermochemistry Vocabulary**

Energy

Matter

Heat

Work

System

Surroundings

Open System

Closed System

Isothermal System

Adiabatic System

Exothermic

Endothermic

**Heat and Work Quantified**

Units

1 calorie = 4.184 joules

1000 calories = 1 Calorie

101.325 Joules = 1 Liter atm

**Heat**

Heat Capacity

Specific Heat

**Work****Energy**

We talk about two types:

Kinetic Energy

Potential Energy

**State Function**

1) What is the heat involved when 250.0 grams of water are heated from 25°C to 75°C?

2) What is the temperature change when 75.0 g of Aluminum (heat capacity 0.900 joule/gram degree) absorbs 250 joules of heat?

3) What is the work when a gas is expanded from 10.0 to 100.0 liters at a pressure of 1.50 atm?

4) If a piston does 50.0 joules of work at 2.00 atm applied pressure how much can you expand a gas?

5) A baseball weighing 0.14 kg is dropped from a building top. When it is 25.0 m from the ground it is moving 15m/s. What is its total energy at this point?

6) An empty swimming pool is 3.5 m deep. A kid on a skateboard (total mass 50 kg) rides down the side of the pool. If he moves with no friction what is his potential energy before he starts to ride down? What is his kinetic energy at the bottom assuming 100% energy transfer? What is his speed at the bottom of the pool?


## **The First Law of Thermodynamics**

We use these ideas to develop and define Enthalpy:


What do the silly little circles mean? "Naught"

**Hess' Law-** Heats of reaction are additive.


1) Given:





Find the enthalpy change for the following reaction:




2) Use the following equations to calculate the enthalpy change for the following:






3) Use the following equations to calculate the enthalpy change for the following:





4) Use the following equations to calculate the enthalpy change for the following:



5) Use the following equations to calculate the enthalpy change for the following:



6) Use the following equations to calculate the enthalpy change for the following:



## **Phase Changes**

What are the six phase changes?

What happens on the molecular level when you heat something?

So what is the dilemma when a phase change occurs?

**Latent Heat**

**Heat of Fusion**

**Heat of Vaporization**

**Heat of Sublimation?**

## Heating Curves

Heat Capacities for common materials:

Substance	Specific Heat	Substance	Specific Heat
H <sub>2</sub> O (s)	2.06 J/g °C	Aluminum (s)	0.900 J/g °C
H <sub>2</sub> O (g)	2.02 J/g °C	Benzene (l)	1.74 J/g °C
H <sub>2</sub> O (l)	4.18 J/g °C	Ethanol (l)	2.42 J/g °C

Phase Change Data

Substance	Heats of Fusion (H <sub>f</sub> ):	Heats of Vaporization (H <sub>v</sub> ):	Boiling Points	Melting Points
H <sub>2</sub> O	6.01 kJ/mol	40.7 kJ/mol	373.2 K	273.2 K
Benzene	10.59 kJ/mol	30.8 kJ/mol	353.2 K	278.6 K
Ethanol	4.60 kJ/mol	43.5 kJ/mol	351.5 K	158.7 K
Acetone	5.72 kJ/mol	29.1 kJ/mol	329.4 K	179 K

Some useful data:

Benzene = C<sub>6</sub>H<sub>6</sub>      Ethanol = C<sub>2</sub>H<sub>5</sub>OH      Acetone = CH<sub>3</sub>COCH<sub>3</sub>

1. How much heat is required to melt 50.0 g of ice at 0°C?
  
  
  
  
  
  
  
  
  
  
2. How much heat is required to melt 75.0 g of Benzene at 278.6 K?







3. You have a sample of H<sub>2</sub>O with a mass of 25.0 g at a temperature of  $-30.0\text{ }^{\circ}\text{C}$ . How many kilojoules of heat energy are necessary to:

- a) heat the ice to  $0^{\circ}\text{C}$ ?
- b) melt the ice?
- c) heat the water from  $0^{\circ}\text{C}$  to  $100^{\circ}\text{C}$ ?
- d) boil the water?
- e) heat the steam from  $100^{\circ}\text{C}$  to  $130^{\circ}\text{C}$ ?





4. How much heat is required to raise 250.0 g of ice at a temperature  $-50.0^{\circ}\text{C}$  to  $125.0^{\circ}\text{C}$ ?

**Heat of Formation:**


Write the Heat of Formation thermochemical equations for the following substances:



Calculate the enthalpy change for the following reactions from standard data tables.



5) For the reaction:



The Heat of Reaction is  $-57.24$  kJ/mole. If the Heat of Formation for  $\text{N}_2\text{O}_4(\text{g})$  is  $9.16$  kJ/mole what is the heat of formation for  $\text{NO}_2(\text{g})$  ?

**Calorimetry**

The first law says that energy can not be created or destroyed, therefore any heat going in has to be accounted for going out!

1) A calorimeter contained 60.0 g of water at 20.00° Celsius. A 100.0 g sample of iron at 100.00°C was added giving the mixture a final temperature of 29.68°C. Calculate the heat capacity of the calorimeter. The specific heat of iron is 0.450 J/g °C.

2) You start with 100 grams of cold water (15°C) in a calorimeter whose heat capacity is 50 cal/°C. To it you add some hot water (80°C). The final temperature is 30°C. What mass of hot water did you add?


3) To a calorimeter (whose heat capacity is  $195 \text{ J/}^\circ\text{C}$ ) containing 100. grams of water at  $25.0^\circ\text{C}$  is added 50.0 grams of aluminum (heat capacity of  $0.900 \text{ J/g}^\circ\text{C}$ ) at  $100.0^\circ\text{C}$ . What is the final temperature?

4) Suppose you have a coffee cup calorimeter whose heat capacity is  $50 \text{ J/}^\circ\text{C}$  that holds 200 mL's of water at  $22^\circ\text{C}$ . To this you drop in a 100.0gram chunk of aluminum that has a specific heat of  $0.900 \text{ J/g}^\circ\text{C}$ . If the metal was at  $95^\circ\text{C}$  what was the final temperature of the mixture?

**Bond Energy**

Where is the energy stored in a chemical reaction?

Find the energy in a Carbon-Hydrogen bond in methane CH<sub>4</sub>.



Now the Carbon-Carbon bond in Ethane. The heat of formation of ethane is  $-84.7 \text{ kJ}$ .

Find the bond energy of a carbon to carbon double bond in the molecule  $C_2H_4(g)$ ,  
 $\Delta H^\circ_f = 52.47 \text{ kJ/mole}$ .

Find the bond energy of a carbon to carbon triple bond in the molecule  $C_2H_2(g)$ ,  
 $\Delta H^\circ_f = 227 \text{ kJ/mole}$ .

Find the bond energy carbon oxygen double bond in Formaldehyde,  $HCOH$ .

$\Delta H^\circ_f = -116 \text{ kJ/mole}$

$\frac{1}{2}O_2(g) \rightleftharpoons O(g)$

$\Delta H^\circ = 247.5 \text{ kJ}$


Bond Energies can also be used to estimate the enthalpy change in a reaction using the idea that energy goes in and out:

Bond Energies:


H-H 432 KJ/mol                      Cl-Cl 239 KJ/mol                      F-F 154 KJ/mol

Br-Br 193 KJ/mol                      H-Cl 427 KJ/mol                      H-F 565 KJ/mol

Use bond energy values from the table to calculate the enthalpy change for the following reaction:



Use bond energy values from the table to calculate the enthalpy change for the following reaction:



Using the heat of reaction from the following equation:



Calculate the bond energy for a H-Br bond.